




CELESTIAL MECHANICS

Vector Derivation of Kepler's Laws from Newtonian Gravitation and the Analytic Determination of Orbital Parameters

Academic Research Paper

Theoretical Astrophysics — Step-by-Step Analytic Derivations, Vector Diagrams and Numerical Tables

Vector Derivation of Kepler's Laws from Newtonian Gravitation and the Analytic Determination of Orbital Parameters

Celestial Mechanics

September 2026

Abstract. Newtonian gravitation gives a remarkably complete analytic theory of isolated planetary, stellar, and spacecraft orbits. This paper presents a vector derivation of Keplerian motion from the two-body problem, emphasizing the geometrical constants of motion that make the inverse-square force integrable. After reducing the two-body system to an equivalent one-body problem with reduced mass, we derive conservation of linear momentum, specific angular momentum, mechanical energy, and the Laplace–Runge–Lenz eccentricity vector. These invariants imply planar conic-section trajectories with focus at the attracting mass and orbit equation $r = p/(1 + e \cos \nu)$. The radial effective potential is developed to identify turning points and circular-orbit stability. Kepler's second and third laws, the vis-viva equation, and the six classical orbital elements follow from the same invariants. We connect true, eccentric, and mean anomalies through Kepler's equation and illustrate its Newton–Raphson solution numerically. Applications include Earth–Sun motion, Hohmann transfer orbits, binary-star mass determination, and an additional exoplanet radial-velocity example. We also discuss observational tests, osculating elements, perturbations, tides, radiation reaction, and relativistic apsidal precession, with an appendix collecting useful algebraic details.

Keywords: celestial mechanics; Kepler problem; two-body problem; orbital elements; binary stars; exoplanets; transfer orbits

1. Introduction

Kepler's three laws were distilled empirically from Tycho Brahe's planetary observations and published between 1609 and 1619. They state that planetary orbits are ellipses with the Sun at one focus, that the radius vector sweeps equal areas in equal times, and that the square of the orbital period is proportional to the cube of the semimajor axis. Newton's achievement was to show that these laws follow from a single dynamical principle: an inverse-square gravitational attraction acting between masses [2]. In modern language, the Kepler problem is the integrable two-body problem generated by the central potential $-\mu/r$, where $\mu = G(m_1 + m_2)$.

The value of the Newtonian derivation is not merely historical. It provides the zeroth-order model for planetary ephemerides, spectroscopic and visual binary stars, exoplanet radial velocities, pulsar timing, spacecraft mission design, and the local dynamics of accretion discs. Most modern treatments add small perturbations: non-spherical gravity fields, additional bodies, gas drag, radiation pressure, tidal dissipation, and relativistic corrections [7, 6, 5]. These refinements are best understood after the exact Keplerian structure has been made explicit.

This paper develops that structure by vector methods. The conservation of angular momentum is familiar and explains planarity and Kepler's second law. Less familiar, but equally decisive, is the Laplace–Runge–Lenz vector, also called the eccentricity vector. Its conservation is a special symmetry of the inverse-square law and fixes both the shape and orientation of

the orbit. From these constants one obtains conic sections without solving a second-order differential equation component by component.

The presentation is intended for advanced undergraduate and beginning graduate students, but the derivations are written with sufficient detail for research use. Throughout, bold symbols denote vectors and hatted symbols denote unit vectors. The relative separation is $\mathbf{r} = \mathbf{r}_1 - \mathbf{r}_2$, its magnitude is $r = |\mathbf{r}|$, and $\mathbf{v} = \dot{\mathbf{r}}$. The gravitational parameter is

$$\mu \equiv G(m_1 + m_2). \quad (1)$$

The specific angular momentum, energy, and eccentricity vector will be denoted by $\mathbf{h}$, ϵ , and $\mathbf{e}$, respectively.

2. Physical and Mathematical Framework

2.1. Newtonian two-body dynamics

Let two point masses m_1 and m_2 have inertial positions $\mathbf{r}_1$ and $\mathbf{r}_2$. Newton's law of gravitation gives equal and opposite forces,

$$m_1 \ddot{\mathbf{r}}_1 = -Gm_1 m_2 \frac{\mathbf{r}_1 - \mathbf{r}_2}{|\mathbf{r}_1 - \mathbf{r}_2|^3}, \quad (2)$$

$$m_2 \ddot{\mathbf{r}}_2 = +Gm_1 m_2 \frac{\mathbf{r}_1 - \mathbf{r}_2}{|\mathbf{r}_1 - \mathbf{r}_2|^3}. \quad (3)$$

The center-of-mass coordinate is

$$\mathbf{R} = \frac{m_1 \mathbf{r}_1 + m_2 \mathbf{r}_2}{m_1 + m_2}, \quad (4)$$

and the relative coordinate is

$$\mathbf{r} = \mathbf{r}_1 - \mathbf{r}_2. \quad (5)$$

Adding (2) and (3) gives

$$(m_1 + m_2)\ddot{\mathbf{R}} = \mathbf{0}. \quad (6)$$

Thus the center of mass moves uniformly. Subtracting the accelerations gives

$$\ddot{\mathbf{r}} = -G(m_1 + m_2)\frac{\mathbf{r}}{r^3} = -\mu\frac{\mathbf{r}}{r^3}. \quad (7)$$

Hence all orbital shape information is contained in the single relative equation (7).

Definition 1. The reduced mass of the two-body system is

$$m_{\text{red}} = \frac{m_1 m_2}{m_1 + m_2}. \quad (8)$$

The relative motion may be interpreted as the motion of a particle of mass m_{red} in the potential $-Gm_1 m_2/r$.

To see this, write the kinetic energy in center-of-mass and relative variables:

$$T = \frac{1}{2}m_1\dot{\mathbf{r}}_1^2 + \frac{1}{2}m_2\dot{\mathbf{r}}_2^2, \quad (9)$$

$$T = \frac{1}{2}(m_1 + m_2)\dot{\mathbf{R}}^2 + \frac{1}{2}m_{\text{red}}\dot{\mathbf{r}}^2. \quad (10)$$

The potential energy is

$$U(r) = -\frac{Gm_1 m_2}{r} = -\frac{m_{\text{red}}\mu}{r}. \quad (11)$$

Dividing the relative energy by m_{red} gives the specific energy

$$\epsilon = \frac{1}{2}v^2 - \frac{\mu}{r}. \quad (12)$$

The adjective “specific” means per unit reduced mass. This convention is standard in astrodynamics and stellar dynamics because the orbit geometry depends on μ and initial relative phase-space coordinates, not separately on m_{red} .

2.2. Central forces and immediate constants

The force in (7) is central: it is parallel to $\mathbf{r}$. Taking the cross product of $\mathbf{r}$ with (7) yields

$$\mathbf{r} \times \ddot{\mathbf{r}} = -\mu\frac{\mathbf{r} \times \mathbf{r}}{r^3} = \mathbf{0}. \quad (13)$$

Since

$$\frac{d}{dt}(\mathbf{r} \times \mathbf{v}) = \mathbf{v} \times \mathbf{v} + \mathbf{r} \times \dot{\mathbf{v}}, \quad (14)$$

we obtain

$$\frac{d\mathbf{h}}{dt} = \mathbf{0}, \quad \mathbf{h} \equiv \mathbf{r} \times \mathbf{v}. \quad (15)$$

The vector $\mathbf{h}$ is the specific angular momentum. It is perpendicular to both $\mathbf{r}$ and $\mathbf{v}$ for all time, so the orbit is confined to the fixed plane normal to $\mathbf{h}$.

Energy conservation follows by dotting (7) with $\mathbf{v}$:

$$\mathbf{v} \cdot \dot{\mathbf{v}} = -\mu\frac{\mathbf{v} \cdot \mathbf{r}}{r^3}, \quad (16)$$

$$\frac{d}{dt}\left(\frac{1}{2}v^2\right) = -\mu\frac{\dot{r}}{r^2} = \frac{d}{dt}\left(\frac{\mu}{r}\right). \quad (17)$$

Therefore

$$\frac{d}{dt}\left(\frac{1}{2}v^2 - \frac{\mu}{r}\right) = 0, \quad (18)$$

which is precisely (12).

Remark 1. Conservation of $\mathbf{h}$ follows from rotational symmetry, while conservation of ϵ follows from time-translation symmetry. The inverse-square force has an additional hidden symmetry, expressed by the Laplace–Runge–Lenz vector, which is responsible for closed bound orbits.

2.3. Radial reduction and the effective potential

Because the orbit is planar, it is useful to introduce polar coordinates in the plane perpendicular to $\mathbf{h}$. The squared speed decomposes as

$$v^2 = \dot{r}^2 + r^2\dot{\nu}^2. \quad (19)$$

The angular-momentum integral gives

$$\dot{\nu} = \frac{h}{r^2}. \quad (20)$$

Substituting (20) into (19) and then into (12) gives a one-dimensional radial energy equation,

$$\frac{1}{2}\dot{r}^2 + U_{\text{eff}}(r) = \epsilon, \quad (21)$$

where the effective potential per unit reduced mass is

$$U_{\text{eff}}(r) = \frac{h^2}{2r^2} - \frac{\mu}{r}. \quad (22)$$

The first term in (22) is the centrifugal barrier. It is not a new force, but the energetic consequence of angular momentum: orbits with $h \neq 0$ cannot reach $r = 0$ unless their energy is high enough to overcome the divergent $h^2/(2r^2)$ term.

The radial acceleration follows by differentiating (21) or by resolving (7) into polar components:

$$\ddot{r} - \frac{h^2}{r^3} = -\frac{\mu}{r^2}. \quad (23)$$

A circular orbit has constant radius $r = r_c$, so $\dot{r} = \ddot{r} = 0$. Equation (23) then gives

$$h^2 = \mu r_c. \quad (24)$$

Equivalently, the circular speed satisfies

$$v_c = \sqrt{\frac{\mu}{r_c}}. \quad (25)$$

The second derivative of the effective potential is

$$\frac{d^2 U_{\text{eff}}}{dr^2} = \frac{3h^2}{r^4} - \frac{2\mu}{r^3}. \quad (26)$$

At the circular radius, (24) gives

$$\left.\frac{d^2 U_{\text{eff}}}{dr^2}\right|_{r_c} = \frac{\mu}{r_c^3} > 0. \quad (27)$$




Figure 1. Effective potential for fixed angular momentum, scaled so that the minimum occurs at $r = r_c = h^2/\mu$. Bound radial motion occurs between turning points satisfying $\epsilon = U_{\text{eff}}(r)$; the minimum is the stable circular orbit.

Thus circular Keplerian orbits are radially stable. Small radial oscillations about r_c have frequency

$$\kappa = \sqrt{\frac{\mu}{r_c^3}}, \quad (28)$$

which equals the angular frequency

$$n_c = \frac{v_c}{r_c} = \sqrt{\frac{\mu}{r_c^3}}. \quad (29)$$

The equality $\kappa = n_c$ is another way of seeing why nearly circular Keplerian orbits close after one radial oscillation. In extended mass distributions or relativistic potentials the equality generally fails, and the apsidal line precesses.

3. The Laplace–Runge–Lenz Vector and the Orbit Equation

3.1. Construction of the eccentricity vector

Starting from (7), take the cross product with $\mathbf{h}$:

$$\dot{\mathbf{v}} \times \mathbf{h} = -\mu \frac{\mathbf{r}}{r^3} \times (\mathbf{r} \times \mathbf{v}). \quad (30)$$

Use the triple-product identity $\mathbf{a} \times (\mathbf{b} \times \mathbf{c}) = \mathbf{b}(\mathbf{a} \cdot \mathbf{c}) - \mathbf{c}(\mathbf{a} \cdot \mathbf{b})$ to write

$$\frac{\mathbf{r}}{r^3} \times (\mathbf{r} \times \mathbf{v}) = \frac{1}{r^3} [\mathbf{r}(\mathbf{r} \cdot \mathbf{v}) - \mathbf{v}r^2], \quad (31)$$

$$= \frac{\dot{r}}{r^2} \mathbf{r} - \frac{1}{r} \mathbf{v}. \quad (32)$$

The derivative of the radial unit vector $\hat{\mathbf{r}} = \mathbf{r}/r$ is

$$\frac{d\hat{\mathbf{r}}}{dt} = \frac{\mathbf{v}}{r} - \frac{\mathbf{r}\dot{r}}{r^2}, \quad (33)$$

$$-\frac{d\hat{\mathbf{r}}}{dt} = \frac{\dot{r}}{r^2} \mathbf{r} - \frac{\mathbf{v}}{r}. \quad (34)$$

Combining (30)–(34) gives

$$\dot{\mathbf{v}} \times \mathbf{h} = \mu \frac{d\hat{\mathbf{r}}}{dt}. \quad (35)$$

Because $\dot{\mathbf{h}} = \mathbf{0}$,

$$\frac{d}{dt}(\mathbf{v} \times \mathbf{h}) = \mu \frac{d\hat{\mathbf{r}}}{dt}. \quad (36)$$

Thus the vector

$$\mathbf{e} = \frac{\mathbf{v} \times \mathbf{h}}{\mu} - \hat{\mathbf{r}} \quad (37)$$

is constant.

Definition 2. The constant vector $\mathbf{e}$ in (37) is the eccentricity vector. Its magnitude

$$e = |\mathbf{e}| \quad (38)$$

is the orbital eccentricity, and for non-circular orbits its direction points toward periapsis.

The physical interpretation is immediate. The vector $\hat{\mathbf{r}}$ follows the instantaneous radius, while $(\mathbf{v} \times \mathbf{h})/\mu$ combines velocity and angular momentum. Their difference remains fixed only for the inverse-square law. This fixed vector chooses a preferred direction in the orbital plane, breaking the rotational degeneracy of a generic central orbit and making the apsidal line stationary.

3.2. Derivation of the conic equation

Dot (37) with $\mathbf{r}$:

$$\mathbf{e} \cdot \mathbf{r} = \frac{\mathbf{r} \cdot (\mathbf{v} \times \mathbf{h})}{\mu} - r. \quad (39)$$

The scalar triple product may be cycled:

$$\mathbf{r} \cdot (\mathbf{v} \times \mathbf{h}) = \mathbf{h} \cdot (\mathbf{r} \times \mathbf{v}) = \mathbf{h} \cdot \mathbf{h} = h^2. \quad (40)$$

Let ν be the true anomaly, the angle from $\mathbf{e}$ to $\mathbf{r}$ in the direction of motion. Then

$$\mathbf{e} \cdot \mathbf{r} = er \cos \nu. \quad (41)$$

Substitution into (39) gives

$$er \cos \nu = \frac{h^2}{\mu} - r. \quad (42)$$

Solving for r yields

$$r = \frac{p}{1 + e \cos \nu}, \quad p \equiv \frac{h^2}{\mu}. \quad (43)$$

Equation (43) is the polar equation of a conic section with one focus at the origin.

Proposition 1. The relative orbit under Newtonian gravity is a conic section. It is an ellipse for $0 \leq e < 1$, a parabola for $e = 1$, and a hyperbola for $e > 1$.

Proof. The polar equation of a conic with focus at the origin, eccentricity e , and semi-latus rectum p is exactly (43). The standard classification of conics by eccentricity gives the three cases. $\square$

At periapsis $\nu = 0$, and at apoapsis for an ellipse $\nu = \pi$. Therefore

$$r_p = \frac{p}{1+e}, \quad (44)$$

$$r_a = \frac{p}{1-e} \quad (e < 1). \quad (45)$$

For an ellipse,

$$a = \frac{r_p + r_a}{2}, \quad (46)$$

$$p = a(1 - e^2). \quad (47)$$

Combining (47) with $p = h^2/\mu$ gives

$$h^2 = \mu a(1 - e^2). \quad (48)$$

For a hyperbola one often uses $a < 0$, in which case the identity

$$p = a(1 - e^2) \quad (49)$$

still gives $p > 0$ because $e > 1$.

The radial turning points of an ellipse may also be recovered from the effective-potential equation. Setting $\dot{r} = 0$ in (21) gives

$$\epsilon r^2 + \mu r - \frac{h^2}{2} = 0. \quad (50)$$

For $\epsilon < 0$ the two positive roots are

$$r_{\pm} = \frac{-\mu \mp \sqrt{\mu^2 + 2\epsilon h^2}}{2\epsilon}. \quad (51)$$

Using (71) below, these roots reduce to

$$r_- = \frac{h^2/\mu}{1+e} = r_p, \quad (52)$$

$$r_+ = \frac{h^2/\mu}{1-e} = r_a. \quad (53)$$

Thus the geometrical apsides are the same points at which the radial kinetic energy vanishes.

4. Kepler's Laws from the Constants of Motion

4.1. Kepler's second law

The areal velocity is the rate at which the radius vector sweeps area in the orbital plane. During an interval dt , the swept area is

$$dA = \frac{1}{2}r^2 dv. \quad (54)$$

In polar coordinates in the orbital plane,

$$\boldsymbol{v} = \dot{r} \hat{\boldsymbol{r}} + r \dot{\nu} \hat{\boldsymbol{\nu}}. \quad (55)$$

Therefore

$$h = |\mathbf{r} \times \mathbf{v}| = r^2 \dot{\nu}. \quad (56)$$




Figure 2. Ellipse geometry. The attracting body is at focus F ; the true anomaly ν is measured at the focus, while the eccentric anomaly E is measured at the center using the auxiliary circle.




Figure 3. Family of Keplerian conics in polar form $r/p = (1 + e \cos \nu)^{-1}$. Ellipses close for $e < 1$; the hyperbolic branch is open and has asymptotes where the denominator vanishes.

Combining (54) and (56),

$$\frac{dA}{dt} = \frac{h}{2}. \quad (57)$$

Since h is constant, dA/dt is constant. This is Kepler's second law.

The physical content is angular-momentum conservation. A body moves rapidly near periapsis because r is small, and slowly near apoapsis because r is large, in exactly the proportion required to sweep equal areas in equal times.

4.2. Kepler's first law

For bound motion $e < 1$, (43) is an ellipse. Its focus is the origin of the relative coordinate, i.e. the center of force. In the exact two-body problem neither star or planet sits at the absolute origin; rather, both bodies orbit their common center of mass. Their relative ellipse

has semimajor axis $a = a_1 + a_2$, where

$$a_1 = \frac{m_2}{m_1 + m_2} a, \quad (58)$$

$$a_2 = \frac{m_1}{m_1 + m_2} a. \quad (59)$$

Thus each body traces a similar ellipse about the barycenter, with the same period and eccentricity but with semimajor axes inversely proportional to mass.

4.3. Kepler's third law

For an ellipse the total area is

$$A_{\text{ell}} = \pi ab. \quad (60)$$

The semiminor axis satisfies

$$b = a\sqrt{1 - e^2}. \quad (61)$$

Since the areal velocity is $h/2$, the period is

$$P = \frac{A_{\text{ell}}}{h/2} = \frac{2\pi a^2 \sqrt{1 - e^2}}{h}. \quad (62)$$

Using (48),

$$P = \frac{2\pi a^2 \sqrt{1 - e^2}}{\sqrt{\mu a(1 - e^2)}}, \quad (63)$$

$$P = 2\pi \sqrt{\frac{a^3}{\mu}}. \quad (64)$$

Therefore

$$P^2 = \frac{4\pi^2}{G(m_1 + m_2)} a^3. \quad (65)$$

This is the Newtonian form of Kepler's third law. The proportionality constant contains the total mass, not merely the central mass. The approximation $P^2 \simeq 4\pi^2 a^3 / (Gm_1)$ is valid only when $m_2 \ll m_1$.

5. Energy, Vis-Viva, and Orbit Classification

5.1. Energy in terms of orbital elements

The eccentricity vector supplies a useful relation between energy and eccentricity. From (37),

$$\mu e = \mathbf{v} \times \mathbf{h} - \mu \hat{\mathbf{r}}. \quad (66)$$

Taking the squared magnitude,

$$\mu^2 e^2 = |\mathbf{v} \times \mathbf{h}|^2 - 2\mu \hat{\mathbf{r}} \cdot (\mathbf{v} \times \mathbf{h}) + \mu^2. \quad (67)$$

Because $\mathbf{v}$ is perpendicular to $\mathbf{h}$,

$$|\mathbf{v} \times \mathbf{h}|^2 = v^2 h^2. \quad (68)$$

Also,

$$\hat{\mathbf{r}} \cdot (\mathbf{v} \times \mathbf{h}) = \frac{1}{r} \mathbf{r} \cdot (\mathbf{v} \times \mathbf{h}) = \frac{h^2}{r}. \quad (69)$$

Thus

$$e^2 = 1 + \frac{h^2}{\mu^2} \left(v^2 - \frac{2\mu}{r} \right). \quad (70)$$

Using $\epsilon = v^2/2 - \mu/r$,

$$e^2 = 1 + \frac{2\epsilon h^2}{\mu^2}. \quad (71)$$

This single equation classifies the orbit:

$$\epsilon < 0 \iff 0 \leq e < 1, \quad (72)$$

$$\epsilon = 0 \iff e = 1, \quad (73)$$

$$\epsilon > 0 \iff e > 1. \quad (74)$$

For an ellipse, substitute (48) into (71):

$$e^2 = 1 + \frac{2\epsilon}{\mu^2} \mu a(1 - e^2), \quad (75)$$

$$1 = -\frac{2\epsilon a}{\mu}. \quad (76)$$

Hence

$$\epsilon = -\frac{\mu}{2a}. \quad (77)$$

The same formula holds for hyperbolic motion if $a < 0$, giving $\epsilon > 0$.

5.2. The vis-viva equation

Insert (77) into the definition of specific energy:

$$\frac{1}{2} v^2 - \frac{\mu}{r} = -\frac{\mu}{2a}. \quad (78)$$

Solving for v^2 gives the vis-viva equation

$$v^2 = \mu \left(\frac{2}{r} - \frac{1}{a} \right). \quad (79)$$

At periapsis and apoapsis,

$$v_p = \sqrt{\mu \frac{1 + e}{a(1 - e)}}, \quad (80)$$

$$v_a = \sqrt{\mu \frac{1 - e}{a(1 + e)}}. \quad (81)$$

Equation (79) is one of the most useful results in celestial mechanics. It says that speed is determined locally by r and globally by the energy scale a . It is therefore central in orbit determination and mission design.

5.3. Flight-path angle and local velocity components

The speed alone does not specify the direction of motion. The radial and transverse components are obtained by differentiating (43) and using (56). Since

$$\frac{dr}{dv} = \frac{pe \sin v}{(1 + e \cos v)^2}, \quad (82)$$

the radial velocity is

$$\dot{r} = \frac{dr}{dv} \dot{v} = \frac{\mu}{h} e \sin v. \quad (83)$$

The transverse velocity is

$$r\dot{v} = \frac{h}{r} = \frac{\mu}{h}(1 + e \cos v). \quad (84)$$

Thus

$$v^2 = \left(\frac{\mu}{h}\right)^2 [e^2 \sin^2 v + (1 + e \cos v)^2], \quad (85)$$

which reduces to the vis-viva form after substituting $p = h^2/\mu$. The flight-path angle γ , measured from the local transverse direction toward the velocity vector, satisfies

$$\tan \gamma = \frac{\dot{r}}{r\dot{v}} = \frac{e \sin v}{1 + e \cos v}. \quad (86)$$

This relation is routinely used in trajectory design because it gives the direction of the required velocity vector at any point on the conic.

6. Classical Orbital Elements

6.1. Geometrical definition

A Keplerian orbit in three-dimensional inertial space is specified by six classical elements:

$$(a, e, i, \Omega, \omega, v). \quad (87)$$

Here a is the semimajor axis, e the eccentricity, i the inclination of the orbital plane relative to a reference plane, Ω the longitude of the ascending node, ω the argument of periapsis, and v the true anomaly.

Let $\hat{k}$ be the unit vector normal to the reference plane. The node vector is

$$\mathbf{n} = \hat{k} \times \mathbf{h}. \quad (88)$$

Then

$$i = \arccos\left(\frac{h_z}{h}\right), \quad (89)$$

$$\Omega = \text{atan2}(n_y, n_x), \quad (90)$$

$$\omega = \text{atan2}\left(\frac{\mathbf{h} \cdot (\mathbf{n} \times \mathbf{e})}{hne}, \frac{\mathbf{n} \cdot \mathbf{e}}{ne}\right), \quad (91)$$

$$v = \text{atan2}\left(\frac{\mathbf{h} \cdot (\mathbf{e} \times \mathbf{r})}{her}, \frac{\mathbf{e} \cdot \mathbf{r}}{er}\right). \quad (92)$$

The two-argument arctangent places the angle in the correct quadrant. For circular or equatorial orbits some angles become undefined; nonsingular element sets are then preferable.

The inverse transformation begins in the perifocal frame $(\hat{p}, \hat{q}, \hat{w})$, where $\hat{p}$ points to periapsis and $\hat{w}$ is normal to the orbit. In that frame,

$$\mathbf{r}_{\text{pf}} = \frac{p}{1 + e \cos v} \begin{pmatrix} \cos v \\ \sin v \\ 0 \end{pmatrix}, \quad (93)$$

$$\mathbf{v}_{\text{pf}} = \sqrt{\frac{\mu}{p}} \begin{pmatrix} -\sin v \\ e + \cos v \\ 0 \end{pmatrix}. \quad (94)$$




Figure 4. Classical orbital elements in a three-dimensional reference frame. The angles Ω , i , and ω orient the orbital plane and apsidal line; the true anomaly v locates the body on the conic.

The inertial vector is obtained by rotations:

$$\mathbf{r} = R_3(\Omega)R_1(i)R_3(\omega)\mathbf{r}_{\text{pf}}, \quad (95)$$

and similarly for $\mathbf{v}$:

$$\mathbf{v} = R_3(\Omega)R_1(i)R_3(\omega)\mathbf{v}_{\text{pf}}. \quad (96)$$

6.2. Anomalies and Kepler's equation

For elliptic motion define the eccentric anomaly E by the auxiliary circle in Fig. 2. The Cartesian coordinates relative to the ellipse center are

$$x_c = a \cos E, \quad (97)$$

$$y_c = b \sin E. \quad (98)$$

With the focus as origin,

$$r = a(1 - e \cos E). \quad (99)$$

Comparison with $r = a(1 - e^2)/(1 + e \cos v)$ yields

$$\cos v = \frac{\cos E - e}{1 - e \cos E}, \quad (100)$$

$$\sin v = \frac{\sqrt{1 - e^2} \sin E}{1 - e \cos E}. \quad (101)$$

Equivalently,

$$\tan \frac{v}{2} = \sqrt{\frac{1+e}{1-e}} \tan \frac{E}{2}. \quad (102)$$

The mean anomaly M is proportional to time from periapsis:

$$M = n(t - \tau), \quad n = \frac{2\pi}{P} = \sqrt{\frac{\mu}{a^3}}, \quad (103)$$

where τ is the time of periapsis passage. Since M is also 2π times the fraction of orbital area swept, one obtains Kepler's equation

$$M = E - e \sin E. \quad (104)$$

A direct derivation follows from area. The area swept from periapsis to eccentric anomaly E is

$$A(E) = \frac{ab}{2}(E - e \sin E). \quad (105)$$

Dividing by πab and multiplying by 2π gives

$$\frac{2\pi A(E)}{\pi ab} = E - e \sin E = M. \quad (106)$$

Thus uniform motion in mean anomaly maps nonlinearly to physical position.

For hyperbolic trajectories, a useful analogue is the hyperbolic anomaly H . With $a < 0$ and $e > 1$,

$$r = a(1 - e \cosh H), \quad (107)$$

where the right-hand side is positive because $a < 0$. The hyperbolic Kepler equation is

$$M_h = e \sinh H - H, \quad (108)$$

with

$$M_h = \sqrt{\frac{\mu}{(-a)^3}}(t - \tau). \quad (109)$$

The true anomaly is related to H by

$$\tan \frac{\nu}{2} = \sqrt{\frac{e+1}{e-1}} \tanh \frac{H}{2}. \quad (110)$$

These formulae are essential for cometary flybys and gravity-assist trajectories, where the object is not gravitationally bound to the central body.

6.3. Newton–Raphson solution

For given M and e , Kepler's equation is solved by finding the root of

$$f(E) = E - e \sin E - M. \quad (111)$$

Newton–Raphson iteration gives

$$E_{j+1} = E_j - \frac{E_j - e \sin E_j - M}{1 - e \cos E_j}. \quad (112)$$

For $0 \leq e < 1$, the derivative satisfies

$$1 - e \leq f'(E) = 1 - e \cos E \leq 1 + e. \quad (113)$$

Thus f is strictly increasing and the root is unique modulo 2π . Once the iteration is sufficiently close, the error is quadratic:

$$|E_{j+1} - E_*| \simeq \left| \frac{f''(E_*)}{2f'(E_*)} \right| |E_j - E_*|^2. \quad (114)$$

For small and moderate eccentricities, $E_0 = M$ is usually adequate. For $e \rightarrow 1$ and M near zero, better starters are used because $f'(E)$ can become small.

Table 1. Newton–Raphson solution of $E - 0.7 \sin E = 2.0$. The reference root is $E_* = 2.447683214$ rad.

j	E_j (rad)	$ E_j - E_* $ (rad)	error ratio
0	2.000000000	4.4768×10^{-1}	–
1	2.492929991	4.5247×10^{-2}	1.01×10^{-1}
2	2.448005625	3.2241×10^{-4}	7.13×10^{-3}
3	2.447683231	1.70×10^{-8}	5.27×10^{-5}
4	2.447683214	1.0×10^{-15}	5.9×10^{-8}




Figure 5. Newton–Raphson convergence for the data in Table 1. The nearly vertical drop after the second iteration illustrates quadratic convergence.

7. Worked Numerical Applications

7.1. Earth–Sun orbit

Take

$$GM_\odot = 1.3271244 \times 10^{20} \text{ m}^3 \text{ s}^{-2}, \quad (115)$$

$$a = 1.4959787 \times 10^{11} \text{ m}, \quad (116)$$

$$e = 0.0167. \quad (117)$$

The period from (64) is

$$P = 2\pi \sqrt{\frac{a^3}{GM_\odot}} = 3.1558 \times 10^7 \text{ s} = 365.26 \text{ d}. \quad (118)$$

The perihelion and aphelion distances are

$$r_p = a(1 - e) = 1.4710 \times 10^{11} \text{ m}, \quad (119)$$

$$r_a = a(1 + e) = 1.5210 \times 10^{11} \text{ m}. \quad (120)$$

The corresponding speeds from (80)–(81) are

$$v_p = 3.03 \times 10^4 \text{ m s}^{-1}, \quad (121)$$

$$v_a = 2.93 \times 10^4 \text{ m s}^{-1}. \quad (122)$$

The fractional speed variation is only a few percent, but it is essential for equal-area motion. The same calculation underlies the barycentric correction applied to high-precision stellar radial velocities.

7.2. Hohmann transfer between circular orbits

A Hohmann transfer is the minimum-impulse coplanar two-burn transfer between two circular orbits when impulses occur tangentially at apsides. Let the initial and

final circular radii be $r_1 < r_2$. The transfer ellipse has

$$a_t = \frac{r_1 + r_2}{2}. \quad (123)$$

The circular speeds are

$$v_1 = \sqrt{\frac{\mu}{r_1}}, \quad (124)$$

$$v_2 = \sqrt{\frac{\mu}{r_2}}. \quad (125)$$

The transfer speeds at periapsis and apoapsis follow from vis-viva:

$$v_{t1} = \sqrt{\mu \left(\frac{2}{r_1} - \frac{1}{a_t} \right)}, \quad (126)$$

$$v_{t2} = \sqrt{\mu \left(\frac{2}{r_2} - \frac{1}{a_t} \right)}. \quad (127)$$

The required impulses are

$$\Delta v_1 = v_{t1} - v_1, \quad (128)$$

$$\Delta v_2 = v_2 - v_{t2}, \quad (129)$$

$$\Delta v_{\text{tot}} = \Delta v_1 + \Delta v_2. \quad (130)$$

The transfer time is half the transfer period:

$$t_H = \pi \sqrt{\frac{a_t^3}{\mu}}. \quad (131)$$

For a low Earth orbit with $r_1 = 6678$ km and a geostationary radius $r_2 = 42164$ km, using $\mu_\oplus = 3.986004 \times 10^5 \text{ km}^3 \text{ s}^{-2}$, one finds

$$a_t = 24421 \text{ km}, \quad (132)$$

$$v_1 = 7.73 \text{ km s}^{-1}, \quad (133)$$

$$v_2 = 3.07 \text{ km s}^{-1}, \quad (134)$$

$$v_{t1} = 10.15 \text{ km s}^{-1}, \quad (135)$$

$$v_{t2} = 1.61 \text{ km s}^{-1}. \quad (136)$$

Thus

$$\Delta v_1 = 2.42 \text{ km s}^{-1}, \quad (137)$$

$$\Delta v_2 = 1.46 \text{ km s}^{-1}, \quad (138)$$

$$\Delta v_{\text{tot}} = 3.88 \text{ km s}^{-1}, \quad (139)$$

$$t_H = 1.91 \times 10^4 \text{ s} = 5.31 \text{ h}. \quad (140)$$

These values omit launch losses, finite-burn effects, inclination changes, and atmospheric drag, but they show how the conic solution becomes a practical engineering tool.

7.3. Binary star mass determination

For a visual binary with angular semimajor axis α in arcseconds, distance d in parsecs, and period P in years, the physical semimajor axis in astronomical units is

$$a(\text{AU}) = \alpha(\text{arcsec}) d(\text{pc}). \quad (141)$$




Figure 6. Hohmann transfer between two circular coplanar orbits. The transfer ellipse is tangent to the inner orbit at periapsis and to the outer orbit at apoapsis.

Using units in which $G = 4\pi^2 \text{ AU}^3 \text{ yr}^{-2} M_\odot^{-1}$, Kepler's third law becomes

$$\frac{m_1 + m_2}{M_\odot} = \frac{a^3(\text{AU})}{P^2(\text{yr})}. \quad (142)$$

Suppose $\alpha = 0.50''$, $d = 20.0 \text{ pc}$, and $P = 25.0 \text{ yr}$. Then

$$a = 10.0 \text{ AU}, \quad (143)$$

$$m_1 + m_2 = \frac{10.0^3}{25.0^2} M_\odot = 1.60 M_\odot. \quad (144)$$

If the barycentric semimajor axes are measured separately and satisfy $a_1/a_2 = m_2/m_1$, then the individual masses follow. For example, if $a_1/a_2 = 0.60$, then

$$m_1 = \frac{1}{1 + 0.60} (1.60 M_\odot) = 1.00 M_\odot, \quad (145)$$

$$m_2 = 0.60 M_\odot. \quad (146)$$

This is the fundamental route by which stellar masses are measured. It anchors the empirical mass–luminosity relation and the calibration of stellar evolution models [8, 9].

7.4. Exoplanet radial-velocity signal

The same barycentric motion is observed in exoplanet searches as a periodic Doppler shift of stellar absorption lines [15]. For a planet of mass m_p orbiting a star of mass $m_\star$, the stellar semimajor axis is

$$a_\star = \frac{m_p}{m_\star + m_p} a. \quad (147)$$

The observed line-of-sight velocity is the projection of the stellar orbital velocity. Its semi-amplitude is

$$K = \frac{2\pi a_\star \sin i}{P \sqrt{1 - e^2}}, \quad (148)$$

where the factor $(1 - e^2)^{-1/2}$ accounts for the nonuniform velocity on an eccentric ellipse. Eliminating a with Kepler's third law gives

$$K = \left(\frac{2\pi G}{P} \right)^{1/3} \frac{m_p \sin i}{(m_\star + m_p)^{2/3}} \frac{1}{\sqrt{1 - e^2}}. \quad (149)$$

When $m_p \ll m_\star$, this becomes the convenient scaling

$$K = 28.4 \text{ m s}^{-1} \left(\frac{m_p \sin i}{M_J} \right) \left(\frac{P}{1 \text{ yr}} \right)^{-1/3} \times \left(\frac{m_\star}{M_\odot} \right)^{-2/3} \frac{1}{\sqrt{1 - e^2}}. \quad (150)$$

Here M_J denotes Jupiter's mass. As a worked example, take a hot Jupiter with

$$P = 3.5247 \text{ d}, \quad (151)$$

$$m_\star = 1.00 M_\odot, \quad (152)$$

$$e = 0, \quad (153)$$

$$m_p \sin i = 0.69 M_J. \quad (154)$$

Since $P = 9.650 \times 10^{-3} \text{ yr}$, (150) gives

$$K = 28.4(0.69)(9.650 \times 10^{-3})^{-1/3} = 92 \text{ m s}^{-1}. \quad (155)$$

This amplitude is easily detectable with precision spectrographs and explains why short-period giant planets were the first exoplanets discovered by the radial-velocity method.

The same orbit has semimajor axis

$$a = \left(\frac{G m_\star P^2}{4\pi^2} \right)^{1/3} = 7.73 \times 10^9 \text{ m} = 0.0517 \text{ AU}. \quad (156)$$

If the stellar radius is $R_\star = 1.0 R_\odot$, the approximate geometric transit probability for a circular orbit is

$$\mathcal{P}_{\text{tr}} \simeq \frac{R_\star}{a} = \frac{6.96 \times 10^8}{7.73 \times 10^9} = 0.090. \quad (157)$$

Thus about 9% of randomly oriented systems with this semimajor axis would transit. Keplerian geometry therefore controls both the Doppler amplitude and the probability that photometry will see an eclipse.

8. Extensions, Generalisations, and Observational Tests

8.1. Osculating elements

When a perturbing acceleration is present, the instantaneous state $(\mathbf{r}, \mathbf{v})$ can still be mapped to a Keplerian conic using the same formulae for ϵ , $\mathbf{h}$, and e . The resulting elements are called osculating elements because the conic is tangent to the true path at that instant. If

$$\ddot{\mathbf{r}} = -\mu \frac{\mathbf{r}}{r^3} + \mathbf{a}_{\text{pert}}, \quad (158)$$




Figure 7. Radial-velocity semi-amplitude for circular edge-on planets around a solar-mass star. The Keplerian scaling $K \propto m_p P^{-1/3}$ favors the discovery of massive short-period planets.

then the angular momentum is no longer exactly constant:

$$\frac{d\mathbf{h}}{dt} = \mathbf{r} \times \mathbf{a}_{\text{pert}}. \quad (159)$$

The energy changes at the rate

$$\frac{d\epsilon}{dt} = \mathbf{v} \cdot \mathbf{a}_{\text{pert}}. \quad (160)$$

The eccentricity vector evolves according to

$$\frac{d\mathbf{e}}{dt} = \frac{\mathbf{a}_{\text{pert}} \times \mathbf{h} + \mathbf{v} \times (\mathbf{r} \times \mathbf{a}_{\text{pert}})}{\mu}. \quad (161)$$

Equations (159)–(161) are compact vector forms of the perturbation problem. They show immediately which physical effects alter the size, shape, or orientation of an orbit. A purely transverse thrust changes both energy and angular momentum efficiently, whereas a purely radial perturbation changes the eccentricity vector without exerting torque.

8.2. Projection tests in binary and planetary systems

Keplerian dynamics makes several falsifiable predictions. In a double-lined spectroscopic binary, the ratio of radial-velocity amplitudes is the inverse mass ratio,

$$\frac{K_1}{K_2} = \frac{m_2}{m_1}. \quad (162)$$

The projected semimajor axes are

$$a_1 \sin i = \frac{K_1 P}{2\pi} \sqrt{1 - e^2}, \quad (163)$$

$$a_2 \sin i = \frac{K_2 P}{2\pi} \sqrt{1 - e^2}. \quad (164)$$

Adding the two and using Kepler's third law gives

$$(m_1 + m_2) \sin^3 i = \frac{P(K_1 + K_2)^3}{2\pi G} (1 - e^2)^{3/2}. \quad (165)$$

If eclipses fix $i \simeq 90^\circ$, the masses become direct dynamical measurements rather than model-dependent estimates.

For transiting exoplanets, the stellar mean density can be inferred from the scaled semimajor axis $a/R_\star$ measured in the light curve. For a circular orbit with $m_p \ll m_\star$,

$$\rho_\star = \frac{3\pi}{GP^2} \left(\frac{a}{R_\star} \right)^3. \quad (166)$$

This relation is a powerful consistency check: the density inferred from the transit geometry must agree with stellar models and spectroscopy, unless the orbit is eccentric or additional bodies perturb the timings.

9. Discussion

9.1. Assumptions and limits of validity

The derivation rests on a short list of idealizations. The bodies are treated as point masses, gravity is Newtonian, no other masses exert forces, and the interaction is instantaneous and conservative. Under these assumptions the two-body problem is exactly integrable. Real astronomical systems are not exactly isolated. Planet-planet interactions produce secular precession and resonances; extended bodies have quadrupole and higher multipole moments; close binaries experience tides and mass transfer; small bodies suffer radiation pressure, Poynting–Robertson drag, and gas drag; compact binaries emit gravitational waves [11].

Perturbations are often organized by writing the acceleration as

$$\ddot{\mathbf{r}} = -\mu \frac{\mathbf{r}}{r^3} + \mathbf{a}_{\text{pert}}. \quad (167)$$

The osculating elements are then the Keplerian elements of the instantaneous conic tangent to the true trajectory. If $\mathbf{a}_{\text{pert}}$ is small, the elements evolve slowly compared with the orbital period. The Kepler problem is therefore not discarded; it becomes the moving basis for perturbation theory.

For a third body of mass m_3 at position $\mathbf{r}_3$ relative to the primary, the leading tidal perturbation on a nearby test particle with $r \ll r_3$ has order

$$a_{\text{tide}} \sim Gm_3 \frac{r}{r_3^3}. \quad (168)$$

Relative to the central acceleration μ/r^2 , the small parameter is

$$\eta_{\text{tide}} \sim \frac{m_3}{m_1} \left(\frac{r}{r_3} \right)^3. \quad (169)$$

This dimensionless estimate explains why Keplerian ellipses are excellent for planet–Sun motion over one orbit, yet long-term integrations are required for Solar-System stability.

9.2. Relativistic apsidal precession

General relativity modifies the Schwarzschild two-body problem by adding a small periastron advance. To leading post-Newtonian order the advance per orbit is

$$\Delta\omega_{\text{GR}} = \frac{6\pi G(m_1 + m_2)}{a(1 - e^2)c^2}. \quad (170)$$

For Mercury, using $a = 0.387$ AU and $e = 0.2056$, this gives

$$\Delta\omega_{\text{GR}} = 5.02 \times 10^{-7} \text{ rad orbit}^{-1}. \quad (171)$$

With about 415 orbits per century,

$$\Delta\omega_{\text{GR}} \simeq 43'' \text{ century}^{-1}, \quad (172)$$

the celebrated anomalous precession explained by Einstein's theory [3, 12, 13]. The smallness of (170) in the Solar System reflects the ratio $GM/(ac^2) \ll 1$. Around neutron stars, black holes, and compact white-dwarf binaries, the same correction becomes dynamically important.

9.3. Radiation reaction in compact binaries

For compact binaries the Newtonian ellipse is not stationary because gravitational waves remove energy and angular momentum. Orbit-averaged radiation reaction gives the classic Peters equations [11],

$$\begin{aligned} \frac{da}{dt} &= -\frac{64}{5} \frac{G^3 m_1 m_2 (m_1 + m_2)}{c^5 a^3 (1 - e^2)^{7/2}} \\ &\quad \times \left(1 + \frac{73}{24} e^2 + \frac{37}{96} e^4 \right), \end{aligned} \quad (173)$$

$$\begin{aligned} \frac{de}{dt} &= -\frac{304}{15} e \frac{G^3 m_1 m_2 (m_1 + m_2)}{c^5 a^4 (1 - e^2)^{5/2}} \\ &\quad \times \left(1 + \frac{121}{304} e^2 \right). \end{aligned} \quad (174)$$

Both a and e decrease for an isolated binary. The inspiral is slow when the separation is large but accelerates rapidly as a shrinks because $da/dt \propto a^{-3}$ for small e . This is the dissipative counterpart of the conservative Kepler problem: at each instant the binary is nearly Keplerian, but the constants of motion drift secularly.

9.4. Comparison with observations and modern developments

Keplerian dynamics is directly visible in planetary ephemerides, visual binary orbits, spectroscopic binary radial-velocity curves, and exoplanet transit timing. A single-lined spectroscopic binary yields the mass function

$$f(m) = \frac{PK_1^3(1 - e^2)^{3/2}}{2\pi G} = \frac{m_2^3 \sin^3 i}{(m_1 + m_2)^2}, \quad (175)$$

where K_1 is the radial-velocity semi-amplitude. This relation is simply Kepler's third law plus projection geometry. When eclipses or astrometry give i , individual masses can be obtained.

In galactic dynamics the same inverse-square two-body result is rarely applied without modification because stars move in the collective potential of many bodies [10]. Nevertheless, local encounters, tidal disruption events, wide binaries, and stars orbiting the Galactic-center black hole are all interpreted using Keplerian elements as the leading model. In cosmology and structure formation, gravitational dynamics becomes

many-body and collisionless, but the Newtonian force law remains the subhorizon limit for nonrelativistic matter [16]. The Kepler problem is therefore both a solved problem and a permanent building block.

10. Conclusion

Starting from Newton's inverse-square gravitational law, the two-body problem reduces exactly to the relative motion of a particle with reduced mass in a central potential. Conservation of specific angular momentum fixes the orbital plane and implies Kepler's equal-area law. Conservation of energy gives the vis-viva equation and the relation $\epsilon = -\mu/(2a)$. The effective potential identifies radial turning points and shows the stability of circular orbits. The additional Laplace–Runge–Lenz vector fixes the apsidal direction and yields the conic orbit equation $r = p/(1 + e \cos \nu)$ directly. Bound orbits are ellipses, parabolic trajectories have zero energy, and hyperbolic trajectories have positive energy.

Kepler's third law follows by combining constant areal velocity with the area of an ellipse, giving $P^2 = 4\pi^2 a^3/[G(m_1 + m_2)]$. This total-mass dependence is essential in binary-star astrophysics and exoplanet radial-velocity work. The classical orbital elements provide a compact geometrical description of a Keplerian orbit in space, while Kepler's equation connects uniform time evolution to position along the ellipse. Newton–Raphson iteration solves this equation rapidly for ordinary eccentricities. Hohmann transfers, binary mass measurements, and exoplanet Doppler amplitudes illustrate how the same analytic framework serves spacecraft dynamics, stellar astrophysics, and planetary science. Perturbations and relativity modify, but do not replace, the Keplerian solution: the osculating conic remains the natural reference state for celestial mechanics.

A. Auxiliary Algebra for Elliptic Motion

This appendix collects algebraic steps that are often skipped in compact presentations. Place the focus at the origin and the major axis on the x -axis, with periastron in the positive x direction. The center of the ellipse is displaced by $-ae$ from the focus, so an eccentric-anomaly parametrization gives

$$x = a(\cos E - e), \quad (176)$$

$$y = b \sin E. \quad (177)$$

The radius from the focus is therefore

$$r^2 = a^2(\cos E - e)^2 + b^2 \sin^2 E, \quad (178)$$

$$= a^2 [(\cos E - e)^2 + (1 - e^2) \sin^2 E], \quad (179)$$

$$= a^2(1 - e \cos E)^2. \quad (180)$$

Taking the positive square root gives

$$r = a(1 - e \cos E), \quad (181)$$

which is (99).

The true anomaly satisfies $x = r \cos \nu$ and $y = r \sin \nu$. Equations (176)–(181) imply

$$\cos \nu = \frac{a(\cos E - e)}{a(1 - e \cos E)}, \quad (182)$$

$$\sin \nu = \frac{a\sqrt{1 - e^2} \sin E}{a(1 - e \cos E)}. \quad (183)$$

Dividing (183) by 1 + (182) in the usual half-angle identity gives

$$\tan \frac{\nu}{2} = \frac{\sin \nu}{1 + \cos \nu} = \sqrt{\frac{1 + e}{1 - e}} \tan \frac{E}{2}. \quad (184)$$

The area formula used in deriving Kepler's equation may also be verified directly. The area swept by the focus-to-particle radius is

$$A(E) = \frac{1}{2} \int_0^E \left(x \frac{dy}{dE'} - y \frac{dx}{dE'} \right) dE'. \quad (185)$$

Using

$$\frac{dx}{dE} = -a \sin E, \quad (186)$$

$$\frac{dy}{dE} = b \cos E, \quad (187)$$

the integrand becomes

$$x \frac{dy}{dE} - y \frac{dx}{dE} = ab(1 - e \cos E). \quad (188)$$

Therefore

$$A(E) = \frac{ab}{2} \int_0^E (1 - e \cos E') dE' = \frac{ab}{2} (E - e \sin E), \quad (189)$$

as used in (105). Finally, since the total area is πab , the fraction of the period elapsed is

$$\frac{t - \tau}{P} = \frac{A(E)}{\pi ab} = \frac{E - e \sin E}{2\pi}, \quad (190)$$

which is equivalent to Kepler's equation.

B. Rotation Matrices and State Vectors

The rotations in (95) and (96) may be written explicitly as

$$R_3(\phi) = \begin{pmatrix} \cos \phi & -\sin \phi & 0 \\ \sin \phi & \cos \phi & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad (191)$$

and

$$R_1(i) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos i & -\sin i \\ 0 & \sin i & \cos i \end{pmatrix}. \quad (192)$$

The combined transformation

$$Q = R_3(\Omega) R_1(i) R_3(\omega) \quad (193)$$

has orthonormal columns. Its first two columns are the inertial components of the perifocal unit vectors $\hat{p}$ and $\hat{q}$:

$$\hat{p} = \begin{pmatrix} \cos \Omega \cos \omega - \sin \Omega \sin \omega \cos i \\ \sin \Omega \cos \omega + \cos \Omega \sin \omega \cos i \\ \sin \omega \sin i \end{pmatrix}, \quad (194)$$

$$\hat{q} = \begin{pmatrix} -\cos \Omega \sin \omega - \sin \Omega \cos \omega \cos i \\ -\sin \Omega \sin \omega + \cos \Omega \cos \omega \cos i \\ \cos \omega \sin i \end{pmatrix}. \quad (195)$$

Consequently the inertial position can be written without matrices as

$$\mathbf{r} = r(\cos \nu \hat{p} + \sin \nu \hat{q}), \quad (196)$$

and the inertial velocity as

$$\mathbf{v} = \sqrt{\frac{\mu}{p}}[-\sin \nu \hat{p} + (e + \cos \nu)\hat{q}]. \quad (197)$$

These formulae are often the most convenient implementation of the mapping from orbital elements to Cartesian initial conditions.

References

- [1] J. Kepler, *Astronomia Nova*, Prague, 1609.
- [2] I. Newton, *Philosophiae Naturalis Principia Mathematica*, London, 1687.
- [3] A. Einstein, "Erklärung der Perihelbewegung des Merkur aus der allgemeinen Relativitätstheorie," *Sitzungsberichte der Königlich Preussischen Akademie der Wissenschaften*, 831–839, 1915.
- [4] S. Chandrasekhar, *Principles of Stellar Dynamics*, Dover, New York, 1960.
- [5] H. Goldstein, C. Poole, and J. Safko, *Classical Mechanics*, 3rd ed., Addison-Wesley, San Francisco, 2002.
- [6] J. M. A. Danby, *Fundamentals of Celestial Mechanics*, 2nd ed., Willmann-Bell, Richmond, 1988.
- [7] C. D. Murray and S. F. Dermott, *Solar System Dynamics*, Cambridge University Press, Cambridge, 1999.
- [8] B. W. Carroll and D. A. Ostlie, *An Introduction to Modern Astrophysics*, 2nd ed., Cambridge University Press, Cambridge, 2017.
- [9] R. Kippenhahn, A. Weigert, and A. Weiss, *Stellar Structure and Evolution*, 2nd ed., Springer, Berlin, 2012.
- [10] J. Binney and S. Tremaine, *Galactic Dynamics*, 2nd ed., Princeton University Press, Princeton, 2008.
- [11] P. C. Peters, "Gravitational radiation and the motion of two point masses," *Physical Review*, **136**, B1224–B1232, 1964.
- [12] S. Weinberg, *Gravitation and Cosmology*, Wiley, New York, 1972.
- [13] C. W. Misner, K. S. Thorne, and J. A. Wheeler, *Gravitation*, W. H. Freeman, San Francisco, 1973.
- [14] R. R. Bate, D. D. Mueller, and J. E. White, *Fundamentals of Astrodynamics*, Dover, New York, 1971.
- [15] M. Mayor and D. Queloz, "A Jupiter-mass companion to a solar-type star," *Nature*, **378**, 355–359, 1995.
- [16] Planck Collaboration, "Planck 2018 results. VI. Cosmological parameters," *Astronomy & Astrophysics*, **641**, A6, 2020.